

Aharonov-Bohm experiments in magnetic field without the Landau levels

Miroslav Pardy

Department of Physical Electronics,
and
The Laboratory of the Plasma Physics,
Masaryk University,
Kotlářská 2, 611 37 Brno, Czech Republic
e-mail:pamir@physics.muni.cz

June 17, 2024

Abstract

We discuss the two-slit experiment and the Aharonov-Bohm (AB) experiment in the magnetic field, where an electron produces so called synchrotron radiation. In other words, the photons are emitted from the points of the electron trajectory and it means that the trajectory of electron is visible in the synchrotron radiation spectrum. The axiomatic system of quantum mechanics does not enable to define the trajectory of the elementary particle. The two-slit experiment and the AB experiment in magnetic field are the missing experiments of quantum mechanics. The extension of the discussion to the cosmical rays moving in the magnetic field of the Saturn magnetosphere and its rings is mentioned. It is related to the probe CASSINI.

Key words Two-slit experiment, Aharonov-Bohm experiment, magnetic field, Landau levels.

1 Introduction

The electron moving in the magnetic field produces so called synchrotron radiation from the Landau levels. The production can be described by the classical electrodynamics, or, in the framework of quantum electrodynamics (Pardy, 2007). From the view point of quantum electrodynamics, the photons are emitted from the points of the trajectory of an electron and the trajectory of electron is visible in the synchrotron radiation spectrum. The emission of photons is stochastic and it causes that the trajectory of electrons for instance in accelerators is not stable. The axiomatic system of quantum mechanics does not enable to define the trajectory of the elementary particle, nevertheless the trajectory is physically meaningful. So, the trajectory is in no case the Kantian *Ding an sich*. Mauritsson et al. (2008) used recently the attosecond laser stroboscope to cleanly image electron motion in atom which is the experimental proof of the existence of the electron trajectories.

The visibility of the trajectory in a magnetic field is not the exact analogue of the trajectory of the charged particle in the gas of the Wilson camera, because in this device the track is generated by the charged particle under special conditions. The Wilson cloud chamber is a track detector which is based on the principle that a charged particle passing through any medium ionizes atoms of the medium. The electromagnetic interaction of a charged particle with this medium is strong enough to remove electrons of atoms, or, to ionize the atoms of the medium. The medium of the Wilson chamber is a damp air. If the air is suddenly expanded, then droplets of condensation form. The droplets condensation form around the ions (the Wilson discovery). The droplets may be photographed, revealing the picture of the ionization trail created by the charged particle¹.

Electron moving in the two-slit experiment immersed in the magnetic field radiates photons and it means that the trajectory of electron is visible in the synchrotron spectrum, or, as the track in the CCD camera. Such experiment was never performed and it means that this is the missing experiment of quantum mechanics. The standard quantum mechanics considers only probability of the appearance of a particle and not its trajectory.

The extension of our discussion to the cosmical magnetic field, or, to the Saturn magnetosphere and its rings is mentioned with regard to the probe CASSINI. The solution of the problem in the framework of the hydrodynamical model of quantum mechanics and the nonlinear quantum mechanics is mentioned. In the next section we discuss the two-slit experiment in the magnetic field and then the Aharonov-Bohm effect in the magnetic field. The article is the small modification of the author texts (Pardy, 2008a, 2008b).

¹ Author performed some experiments with the Wilson chamber as a student at the Purkyně University in Brno, Czech Republic

2 The two-slit experiment

An alternative to the standard quantum mechanics, is the de Broglie-Bohm theory where particles also have precise locations at all times, and that their velocities are defined by the wave-function. So, while a single particle will travel through one particular slit in the double-slit experiment, the so-called "pilot wave" that influences it will travel through both. The two slit de Broglie-Bohm trajectories were first calculated by in 1979 (Phillipidis et al., 1979) . The de Broglie-Bohm theory produces the same statistical results as standard quantum mechanics, without many of its conceptual difficulties.

Following Holstein (1992), the electron beam, in the two-slit experiment, is emitted from the left to the right from the source S to the screen S_1 with the upper slit A_1 and the lower slit A_2 . Then it continues to the screen S_2 to form the diffraction pattern. The symbolic scheme is as follows:

$$S \rightarrow S_1 \rightarrow S_2. \quad (1)$$

The corresponding classical trajectories are SA_1P , and SA_2P , P being the point of the impinging electron on the screen S_2 .

Assuming the dominance of the classical trajectories (Holstein, 1992), the phase along the first trajectory is as follows:

$$\exp\left(i \int_{t_1}^{t_2} dt \frac{1}{2} m \dot{\mathbf{x}}^2\right) = \exp\left(i \left[\int_S^{A_1} + \int_{A_1}^P \right] dt \frac{1}{2} mv\right) \approx \exp\left(i \frac{\pi d_0}{\lambda} + i \frac{\pi d_1}{\lambda}\right), \quad (2)$$

where

$$d_0 = SA_1, \quad d_1 = A_1P, \quad \lambda = \frac{2\pi}{p}, \quad (3)$$

where $\lambda = 2\pi/mv$ being the de Broglie wavelength.

The phase along the second trajectory is as follows:

$$\exp\left(i \int_{t_1}^{t_2} dt \frac{1}{2} m \dot{\mathbf{x}}^2\right) \approx \exp\left(i \frac{\pi d_0}{\lambda} + i \frac{\pi d_2}{\lambda}\right); \quad d_2 = A_2P. \quad (4)$$

The relative phase difference between the two paths is then given by the formula

$$\exp\left(i \frac{\pi d_2 - \pi d_1}{\lambda}\right). \quad (5)$$

The wave function at the point on the screen S is then

$$\psi \approx \psi_0 \left(1 + \exp\left[i\pi \frac{d_2 - d_1}{\lambda}\right] \right). \quad (6)$$

The intensity is

$$|\psi|^2 \approx |\psi_0|^2 4 \cos^2 \pi \frac{d_2 - d_1}{2\lambda}. \quad (7)$$

If we denote by L the distance between screens, by δ the distance between slits and by s the distance of the point on the screen S_2 from the axis of symmetry of the experiment, then we can write using the Pythagoras theorem $c^2 = a^2 + b^2$; $c = d_{1,2}$, $a = L$, $b = s \pm \delta/2$:

$$d_2 = \sqrt{L^2 + \left(s + \frac{1}{2}\delta\right)^2} \approx \sqrt{L^2 + s^2} + \frac{s\delta}{2\sqrt{L^2 + s^2}} = \sqrt{L^2 + s^2} + \frac{\delta}{2} \sin \theta, \quad (8)$$

$$d_1 = \sqrt{L^2 + \left(s - \frac{1}{2}\delta\right)^2} \approx \sqrt{L^2 + s^2} - \frac{s\delta}{2\sqrt{L^2 + s^2}} = \sqrt{L^2 + s^2} - \frac{\delta}{2} \sin \theta. \quad (9)$$

So, we can write the intensity distribution of electron in term of the angle θ and the slit separation in the form:

$$I(\theta) = 4I_0 \cos^2 \left(\pi \delta \frac{\sin \theta}{2\lambda} \right), \quad (10)$$

where I_0 is the intensity in case that only single slit is open.

If we switch on the magnetic field between the screen S and S_2 then it is evident that the phase obtained along the first trajectory and the second trajectory is as follows:

$$(phase)_1 \approx \exp \left(i \frac{\pi d_0}{\lambda} + i \frac{\pi d_1}{\lambda} + ie \int_1 \mathbf{A} \cdot d\mathbf{x} \right) \quad (11)$$

and

$$(phase)_2 \approx \exp \left(i \frac{\pi d_0}{\lambda} + i \frac{\pi d_2}{\lambda} + ie \int_2 \mathbf{A} \cdot d\mathbf{x} \right). \quad (12)$$

The phase difference between two trajectories is evidently as follows

$$\Delta(phase) \approx \exp \left(i \frac{\pi d_2}{\lambda} - i \frac{\pi d_1}{\lambda} + ie \oint \mathbf{A} \cdot d\mathbf{x} \right). \quad (13)$$

Using the Stokes theorem

$$ie \oint \mathbf{A} \cdot d\mathbf{x} = ie \int_{area} \text{rot} \mathbf{A} \cdot d\mathbf{S} = ie \int_{area} \mathbf{B} \cdot d\mathbf{S} = ie B S_{area}, \quad (14)$$

where \mathbf{B} is the density of the magnetic induction at the area SA_2PA_1S .

The intensity patterns is then shifted due to the existence of the magnetic field as

$$I(\theta) = 4I_0 \cos^2 \left(\pi \delta \frac{\sin \theta}{2\lambda} + e \frac{1}{2} B S_{area} \right). \quad (15)$$

During the calculation we used the assumption that the magnetic field is sufficiently small to not change the original dominant trajectories and it is very small, to cause

the change of these trajectories by the synchrotron bremsstrahlung. The synchrotron radiation is the crucial effect to see the electron trajectory in reality. Such approach can be used not only in the quantum mechanical two-slit experiments but also in case of detections of the charged particle in the particle laboratories, or, as the synchrotron radiation observation of the cosmical particles when moving in the cosmical magnetic fields. To our knowledge, such experimental approach was not applied till this time.

3 Aharonov-Bohm effect

Werner Ehrenberg and Raymond E. Siday first predicted the effect in 1949 (Ehrenberg et al., 1949). Yakir Aharonov and David Bohm published their analysis in 1959. After publication of the 1959 paper, Bohm was informed of Ehrenberg and Siday's work, which was acknowledged and credited in Bohm and Aharonov's subsequent 1961 paper.

The effect was confirmed experimentally, with a very large error, while Bohm was still alive. By the time the error was down to a respectable value, Bohm had died (Peshkin et al., 1989). The present elaborate is some modification of the original author article (Parady, 2008b).

In case of the so called Bohm-Aharonov effect an infinite solenoid is introduced between slits A_1, A_2 on the right side of the screen S_1 . Since the solenoid is infinite, there is no magnetic field outside the solenoid volume itself. However, There is non-vanishing vector potential outside of the solenoid, which can be expressed in the cylindrical coordinates r, φ, z as

$$\mathbf{A} = \begin{cases} A_\varphi = \frac{1}{2}Br; & r < R, \\ A_\varphi = \frac{1}{2}B\frac{R^2}{r}, & r > R, \end{cases} \quad (16)$$

which corresponds to the magnetic field nonzero inside the solenoid and zero out of the solenoid of the diameter $2R$, or,

$$\mathbf{B} = \text{rot}\mathbf{A} = B_z = \frac{1}{r} \frac{\partial}{\partial r}(rA_\varphi) = \begin{cases} B_z = B; & r < R, \\ B_z = 0 & r > R. \end{cases} \quad (17)$$

The phase difference at the point P between two paths is analogical to the previous discussion

$$\text{phase difference} \approx \exp\left(i\frac{\pi d_2}{\lambda} - i\frac{\pi d_1}{\lambda} + ie \oint \mathbf{A} \cdot d\mathbf{x}\right) \quad (18)$$

where the circle of integration is SA_2PA_1S .

According to the Stokes theorem we get in analogy with eq. (13):

$$ie \oint \mathbf{A} \cdot d\mathbf{x} = ie \int_{area} \text{rot}\mathbf{A} \cdot d\mathbf{S} = ie \int_{area} \mathbf{B} \cdot d\mathbf{S} = ieBS_{area}, \quad (19)$$

The intensity pattern is shifted according in analogy with eq. (14). Or,

$$I(\theta) = 4I_0 \cos^2 \left(\pi\delta \frac{\sin \theta}{2\lambda} + e \frac{1}{2} B \pi R^2 \right). \quad (20)$$

In case that in the Bohm-Aharonov arrangement the external magnetic field is switch on of the intensity B_{ext} , then the total shift of the intensity pattern is obviously:

$$I(\theta) = 4I_0 \cos^2 \left(\pi\delta \frac{\sin \theta}{2\lambda} + e \frac{1}{2} \pi B_{solenoid} R^2 + e \frac{1}{2} B_{ext} S_{area} \right), \quad (21)$$

where in this case S_{area} is the area SA_2PA_1S

We see that the existence of the Landau levels (Landau, 1965) are not present in eq. (21) in our approximation.

While the Bohm-Aharonov effect was verified experimentally, the Bohm-Aharonov effect in the homogeneous magnetic field was, to our knowledge, never performed. Also in this experiment the synchrotron radiation of electron is produced and it means that the trajectories are visible in the synchrotron radiation spectrum. Of course, we can derive the AB effect completely in terms of magnetic field and not in terms of the vector potential. However, with the same result. It is necessary only to stress that the wave function of electron "feels" the magnetic field inside the solenoid and it respects it.

4 Discussion

We have considered the two-slit experiment and the AB experiment in the magnetic field. In such a case the electron moving in the magnetic field produces so called synchrotron radiation. In other words, the photons are emitted from the the points of the trajectory of electron and it means that the trajectory of electron is visible in the synchrotron radiation spectrum. The axiomatic system of quantum mechanics does not enable to define the trajectory of the elementary particle, nevertheless the trajectory is physically meaningful. Electron is an elementary particle with the localized mass and charge and it is geometrically point-like. The two-slit experiment and AB experiment in a magnetic field was never performed and it means that they are the missing experiments of quantum mechanics. The situation can be considered as the analog of the particle detection in the Wilson camera where the track of a charged particle is visible. Trajectories of the particles moving in the magnetic field are visible in the synchrotron radiation spectrum detected by the CCD camera.

Let us remark that the Aharonov-Bohm problem can be reformulated in such a way that the Schrödinger equation is transformed in the form that it uses magnetic intensities instead the vector potential. Then the problem is reduced to the scattering of electrons on the cylindric magnetic field. It means that the Aharonov-Bohm conception is only the theoretical method of calucation of he specific experimental situation and nothing else. Or, according to Weisskopf, physics is only the grand system of methods. Including also the Einstein general theory of relativity and gravity.

There is a dual experiment where the magnetic monopole moves in the electric field and produces the synchrotron radiation. To our knowledge, duality was developed in particle physics but the dual experiments with magnetic monopoles were never performed. So, the dual experiments are also missing ones in quantum electrodynamics.

The analysis can be extended to the cosmical space, where the charged cosmical rays move in the magnetic field and produce the synchrotron radiation spectrum which enables to make the trajectories visible. The opportunity was given to the cosmical probe CASSINI moving on the orbit around the Saturn in its magnetosphere and in the magnetic field of its rings. The charged cosmical rays moving in such magnetic fields generate the synchrotron radiation which can be detected by CCD camera. In other words the trajectories are visible. So, the Auger Argentina cosmic rays project can be supplemented by the Saturn cosmical ray project realized by the probe CASSINI (Matthews et al., 2004).

We frequently read in the physical texts on the quantum mechanics that the classical limit of quantum mechanics is obtained only by the so called WKB method. However, the limit is only formal because in this case the probabilistic form of the solution is conserved while classical mechanics is strongly deterministic. In other words, statistical description of quantum mechanics is in no case reduced to the strong determinism of classical mechanics of one-particle system. So, only nonlinear quantum mechanics of the above form gives the correct classical limit expressed by the delta-function. More information on the problems which are solved by the nonlinear Schrödinger equation involving the collapse of the wave function and the Schrödinger cat paradox is described in author's articles (Pardy, 2001, 1994). The extended version of the nonlinear quantum world is described in the preprint of Castro (2002).

References

- Castro, C., Mahecha, J. and Rodríguez, B., (2002) Nonlinear QM as a fractal Brownian motion with complex diffusion constant, quant-ph/0202026.
- Ehrenberg, W. and Siday, R. E. (1949). "The Refractive Index in Electron Optics and the Principles of Dynamics". Proceedings of the Physical Society B 62, No. 1, 8-21.
- Holstein, B. R., it Topics in advanced quantum mechanics, Addison-Wesley Publishing Company, Redwood City, (1992).
- Landau, L. D. (1992). Diamagnetism of metals, Z. Phys., 04, 629, 1930, in: *Collected papers of L. D. Landau*, Edited and with an introduction by D. T. Haar, (Gordon and Breach Science Publishers, New York, 1965)
- Matthews, L. S. and Hyde, T. W. (2004). Gravitoelectrodynamics in Saturn's F Ring:

Encounters with Prometheus and Pandora, preprint of the Center for Astrophysics, Space Physics, and Engineering Research, Baylor University.

Mauritsson, J. Johnsson, P., Mansten, E., Swoboda, T., Ruchon, A., L'Huillier, A. and Schafer, K. J. (2008). Coherent electron scattering captured by an attosecond quantum stroboscope, *Phys. Rev. Lett.* 100, 073003.

Pardy, M., Possible Tests of Nonlinear Quantum Mechanics, in: *Waves and Particles in Light and Matter*, Ed. by Alwyn van der Merwe and Garuccio, A., Plenum Press New York, (1994).

Pardy, M. (2001). To the nonlinear quantum mechanics, [quant-ph/0111105](#).

Pardy, M. (2007). The synchrotron radiation from the Volkov solution of the Dirac equation: e-print [hep-ph/0703102](#).

Pardy, M. (2008a). Missing experiment in quantum mechanics, [arXiv:0801.2483v1 \[quant-ph\]](#).

Pardy, M. (2008b). Two-Slit and Aharonov-Bohm Experiments in Magnetic Field, *Int. J. Theor. Phys.* (2008) 47: 32733281 DOI 10.1007/s10773-008-9763-5.

Peshkin, M. and Tonomura, A. (1989). Aharonov-Bohm effect, (Springer-Verlag. ISBN 978-3-540-51567-8).

Philippidis, C., Dewdney, C. and Hiley, B. J. (1979). "Quantum interference and the quantum potential". *Il Nuovo Cimento B.* 52, No. 1, 15-28.