

Khovanov Homology

Louis H. Kauffman
Department of Mathematics, Statistics
and Computer Science (m/c 249)
851 South Morgan Street
University of Illinois at Chicago
Chicago, Illinois 60607-7045
<kauffman@uic.edu>

July 11, 2011

Abstract

This paper is an introduction to Khovanov Homology.

Keywords. bracket polynomial, Khovanov homology, cube category, simplicial category, tangle cobordism, chain complex, chain homotopy, unitary transformation, quantum computing, quantum information theory, link homology, categorification.

1 Introduction

This paper is an introduction to Khovanov homology. We start with a quick introduction to the bracket polynomial, reformulating it and the Jones polynomial so that the value of an unknotted loop is $q + q^{-1}$. We then introduce enhanced states for the bracket state sum so that, in terms of these enhanced states the bracket is a sum of monomials. Then in Section 3, we point out that the shape of the collection of bracket states for a given diagram is a cube and that this cube can be taken to be a category. It is an example of a *cube category*. We show that functors from a cube category to a category of modules naturally have homology theories associated with them. In Section 4 we show how to make a homology theory (Khovanov homology) from the states of the bracket so that the enhanced states are the generators of the chain complex. We show how a Frobenius algebra structure arises naturally from this adjacency structure for the enhanced states. Finally we show that the resulting homology is an example of homology related to a module functor on the cube category as described in Section 3. In Section 5, we give a short exposition of Dror BarNatan's tangle cobordism theory for Khovanov homology. This theory replaces Khovanov homology by an abstract chain homotopy class of a complex of surface

cobordisms associated with the states of a knot or link diagram. In Section 6 we give a short exposition of the Rasmussen invariant and its application to finding the four-ball genus of torus knots. In Section 7 we give a description of Khovanov homology as the homology of a simplicial module by following our description of the cube category in this context. I regard this section as a step in the direction of providing an abstract simplicial homotopy theory for Khovanov homology, but this project is not yet completed. In Section 8 we discuss a quantum context for Khovanov homology that is obtained by building a Hilbert space whose orthonormal basis is the set of enhanced states of a diagram K . Then there is a unitary transformation U_K of this Hilbert space so that the Jones polynomial J_K is the trace of U_K : $J_K = \text{Trace}(U_K)$. We discuss a generalization where the linear space of the Khovanov homology itself is taken to be the Hilbert space. In this case we can define a unitary transformation U'_K so that, for values of q and t on the unit circle, the Poincaré polynomial for the Khovanov homology is the trace of U'_K . Section 9 is a discussion of other forms of link homology and categorification with selected references.

It gives the author great pleasure to thank the members of the Quantum Topology Seminar at the University of Illinois at Chicago for many useful conversations and to thank the Perimeter Institute in Waterloo, Canada for their hospitality while this paper was being completed.

2 Bracket Polynomial and Jones Polynomial

The bracket polynomial [15] model for the Jones polynomial [12, 13, 14, 52] is usually described by the expansion

$$\langle \times \rangle = A \langle \smile \rangle + A^{-1} \langle \frown \rangle$$

Here the small diagrams indicate parts of otherwise identical larger knot or link diagrams. The two types of smoothing (local diagram with no crossing) in this formula are said to be of type A (A above) and type B (A^{-1} above).

$$\begin{aligned} \langle \bigcirc \rangle &= -A^2 - A^{-2} \\ \langle K \bigcirc \rangle &= (-A^2 - A^{-2}) \langle K \rangle \\ \langle \diagup \rangle &= (-A^3) \langle \smile \rangle \\ \langle \diagdown \rangle &= (-A^{-3}) \langle \smile \rangle \end{aligned}$$

One uses these equations to normalize the invariant and make a model of the Jones polynomial. In the normalized version we define

$$f_K(A) = (-A^3)^{-wr(K)} \langle K \rangle / \langle \bigcirc \rangle$$

where the writhe $wr(K)$ is the sum of the oriented crossing signs for a choice of orientation of the link K . Since we shall not use oriented links in this paper, we refer the reader to [15] for

the details about the writhe. One then has that $f_K(A)$ is invariant under the Reidemeister moves (again see [15]) and the original Jones polynomial $V_K(t)$ is given by the formula

$$V_K(t) = f_K(t^{-1/4}).$$

The Jones polynomial has been of great interest since its discovery in 1983 due to its relationships with statistical mechanics, due to its ability to often detect the difference between a knot and its mirror image and due to the many open problems and relationships of this invariant with other aspects of low dimensional topology.

The State Summation. In order to obtain a closed formula for the bracket, we now describe it as a state summation. Let K be any unoriented link diagram. Define a *state*, S , of K to be the collection of planar loops resulting from a choice of smoothing for each crossing of K . There are two choices (A and B) for smoothing a given crossing, and thus there are $2^{c(K)}$ states of a diagram with $c(K)$ crossings. In a state we label each smoothing with A or A^{-1} according to the convention indicated by the expansion formula for the bracket. These labels are the *vertex weights* of the state. There are two evaluations related to a state. The first is the product of the vertex weights, denoted $\langle K|S \rangle$. The second is the number of loops in the state S , denoted $\|S\|$. Define the *state summation*, $\langle K \rangle$, by the formula

$$\langle K \rangle = \sum_S \langle K|S \rangle \delta^{\|S\|}$$

where $\delta = -A^2 - A^{-2}$. This is the state expansion of the bracket. It is possible to rewrite this expansion in other ways. For our purposes in this paper it is more convenient to think of the loop evaluation as a sum of *two* loop evaluations, one giving $-A^2$ and one giving $-A^{-2}$. This can be accomplished by letting each state curve carry an extra label of $+1$ or -1 . We describe these *enhanced states* below. But before we do this, it will be useful for the reader to examine Figure 2. In Figure 2 we show all the states for the right-handed trefoil knot, labelling the sites with A or B where B denotes a smoothing that would receive A^{-1} in the state expansion.

Note that in the state enumeration in Figure 2 we have organized the states in tiers so that the state that has only A -smoothings is at the top and the state that has only B -smoothings is at the bottom.

Changing Variables. Letting $c(K)$ denote the number of crossings in the diagram K , if we replace $\langle K \rangle$ by $A^{-c(K)}\langle K \rangle$, and then replace A^2 by $-q^{-1}$, the bracket is then rewritten in the following form:

$$\langle \times \rangle = \langle \smile \rangle - q \langle \rangle \langle \rangle$$

with $\langle \bigcirc \rangle = (q + q^{-1})$. It is useful to use this form of the bracket state sum for the sake of the grading in the Khovanov homology (to be described below). We shall continue to refer to the smoothings labeled q (or A^{-1} in the original bracket formulation) as *B-smoothings*.




Figure 1: **Reidemeister Moves**

We catalog here the resulting behaviour of this modified bracket under the Reidemeister moves.

$$\begin{aligned}
 \langle \bigcirc \rangle &= q + q^{-1} \\
 \langle K \bigcirc \rangle &= (q + q^{-1}) \langle K \rangle \\
 \langle \nearrow \searrow \rangle &= q^{-1} \langle \smile \rangle \\
 \langle \searrow \nearrow \rangle &= -q^2 \langle \smile \rangle \\
 \langle \diagup \diagdown \rangle &= -q \langle \cup \rangle \\
 \langle \diagdown \diagup \rangle &= \langle \diagup \diagdown \rangle
 \end{aligned}$$

It follows that if we define

$$J_K = (-1)^{n_-} q^{n_+ - 2n_-} \langle K \rangle,$$

where n_- denotes the number of negative crossings in K and n_+ denotes the number of positive crossings in K , then J_K is invariant under all three Reidemeister moves. Thus J_K is a version of the Jones polynomial taking the value $q + q^{-1}$ on an unknotted circle.

Using Enhanced States. We now use the convention of *enhanced states* where an enhanced state has a label of 1 or -1 on each of its component loops. We then regard the value of the loop $q + q^{-1}$ as the sum of the value of a circle labeled with a 1 (the value is q) added to the value of a circle labeled with an -1 (the value is q^{-1}). We could have chosen the less neutral labels of $+1$ and X so that

$$q^{+1} \iff +1 \iff 1$$

and

$$q^{-1} \iff -1 \iff x,$$

since an algebra involving 1 and x naturally appears later in relation to Khovanov homology. It does no harm to take this form of labeling from the beginning. The use of enhanced states for formulating Khovanov homology was pointed out by Oleg Viro in [48].

Consider the form of the expansion of this version of the bracket polynomial in enhanced states. We have the formula as a sum over enhanced states s :

$$\langle K \rangle = \sum_s (-1)^{i(s)} q^{j(s)}$$

where $i(s)$ is the number of B -type smoothings in s and $j(s) = i(s) + \lambda(s)$, with $\lambda(s)$ the number of loops labeled 1 minus the number of loops labeled -1 in the enhanced state s .

One advantage of the expression of the bracket polynomial via enhanced states is that it is now a sum of monomials. We shall make use of this property throughout the rest of the paper.

3 Khovanov Homology and the Cube Category

We are going to make a chain complex from the states of the bracket polynomial so that the homology of this chain complex is a knot invariant. One way to see how such a homology theory arises is to step back and note that the collection of states for a diagram K forms a category in the shape of a cube. A functor from such a category to a category of modules gives rise to a homology theory in a natural way, as we explain below.

Examine Figure 2 and Figure 3. In Figure 2 we show all the standard bracket states for the trefoil knot with arrows between them whenever the state at the output of the arrow is obtained from the state at the input of the arrow by a single smoothing of a site of type A to a site of type B . The abstract structure of this collection of states is a category with objects of the form $\langle ABA \rangle$ where this symbol denotes one of the states in the state diagram of Figure 2. In Figure 3 we illustrate this cube category (the states are arranged in the form of a cube) by replacing the states in Figure 2 by symbols $\langle XYZ \rangle$ where each literal is either an A or a B . A typical generating morphism in the 3-cube category is

$$\langle ABA \rangle \longrightarrow \langle BBA \rangle.$$

We formalize this way of looking at the bracket states as follows. Let $\mathcal{S}(K)$ denote a category associated with the states of the bracket for a diagram K whose objects are the states, with sites labeled A and B as in Figure 2. A morphism in this category is an arrow from a state with a given number of A 's to a state with fewer A 's.

Let $\mathcal{D}^n = \{A, B\}^n$ be the n -cube category whose objects are the n -sequences from the set $\{A, B\}$ and whose morphisms are arrows from sequences with greater numbers of A 's to sequences with fewer numbers of A 's. Thus \mathcal{D}^n is equivalent to the poset category of subsets of $\{1, 2, \dots, n\}$. We make a functor $\mathcal{R} : \mathcal{D}^n \longrightarrow \mathcal{S}(K)$ for a diagram K with n crossings as follows. We map sequences in the cube category to bracket states by choosing to label the crossings of the diagram K from the set $\{1, 2, \dots, n\}$, and letting this functor take abstract A 's and B 's in the cube category to smoothings at those crossings of type A or type B . Thus each sequence in the cube category




Figure 2: **Bracket States and Khovanov Complex**




Figure 3: **Cube Category**

is associated with a unique state of K when K has n crossings. By the same token, we define a functor $\mathcal{S} : \mathcal{S}(K) \rightarrow \mathcal{D}^n$ by associating a sequence to each state and morphisms between sequences corresponding to the state smoothings. With these conventions, the two compositions of these morphisms are the identity maps on their respective categories.

Let \mathcal{M} be a pointed category with finite sums, and let $\mathcal{F} : \mathcal{D}^n \rightarrow \mathcal{M}$ be a functor. In our case \mathcal{M} will be a category of modules and \mathcal{F} will carry n -sequences to certain tensor powers corresponding to the standard bracket states of a knot or link K . We postpone this construction for a moment, and point out that there is a natural structure of chain complex associated with the functor \mathcal{F} . First note that each object in \mathcal{D}^n has the form

$$X = \langle X_0 \cdots X_{n-1} \rangle$$

where each X_i equals either A or B and we have morphisms

$$d_i : \langle X_0 \cdots X_i \cdots X_{n-1} \rangle \rightarrow \langle X_0 \cdots \bar{X}_i \cdots X_{n-1} \rangle$$

whenever $X_i = A$ and (by definition) $\bar{X}_i = B$. We then define

$$\partial_i = \mathcal{C}(d_i) : \mathcal{C}\langle X_0 \cdots X_i \cdots X_{n-1} \rangle \rightarrow \mathcal{C}\langle X_0 \cdots \bar{X}_i \cdots X_{n-1} \rangle$$

whenever d_i is defined. We then define the chain complex C by

$$C^k = \bigoplus_X \mathcal{C}\langle X_0 \cdots X_{n-1} \rangle$$

where each sequence $X = \langle X_0 \cdots X_{n-1} \rangle$ has k B 's. With this we define

$$\partial : C^k \rightarrow C^{k+1}$$

by the formula

$$\partial x = \sum_{i=0}^{n-1} (-1)^{c(X,i)} \partial_i(x)$$

for $x \in \mathcal{C}X = \mathcal{C}\langle X_0 \cdots X_{n-1} \rangle$ and $c(X, i)$ denotes the number of A 's in the sequence X that precede X_i .

We want $\partial^2 = 0$ and it is easy to see that this is equivalent to the condition that $\partial_i \partial_j = \partial_j \partial_i$ for $i \neq j$ whenever these maps and compositions are defined. We can assume that the functor \mathcal{F} has this property, or we can build it in axiomatically by adding the corresponding relations to the cube category in the form $d_i d_j = d_j d_i$ for $i \neq j$ whenever these maps are defined. In the next section we shall see that there is a natural way to define the maps in the state category so that this condition holds. Once we axiomatize this commutation relation at the level of the state category or the cube category, then the functor \mathcal{F} will induce a chain complex and homology as above.

In this way, we see that a suitable functor from the cube category to a module category allows us to define homology that is modeled on the ‘‘shape’’ of the cube. The set of bracket states forms a natural functorial image of the cube category, and that makes it possible to define the Khovanov chain complex. In terms of the bracket states, we will map each state loop to a specific module V , and each state to a tensor power of V to the number of loops in the state. The details of this construction are in the next section.

We use a specific construction for the Khovanov complex that is directly related to the enhanced states for the bracket polynomial, as we will see in the next section. In this construction we will use the enhanced states, regarding each loop as labeled with either 1 or x for a module $V = k[x]/(x^2)$ associated with the loop (where $k = \mathbb{Z}/2\mathbb{Z}$ or $k = \mathbb{Z}$.) Thus the two labelings of the loop will correspond to the two generators of the module V . A state that is a collection of loops will be associated with $V^{\otimes r}$ where r is the number of loops in the state. In this way we will obtain a functor from the state category to a module category, and at the same time it will happen that any single enhanced state will correspond to a generator of the chain complex. In the next section we show how naturally this algebra appears in relation to the enhanced states. We then return to the categorical point of view and see how, surface cobordisms of circles provide an abstract category for the invariant.

4 Khovanov Homology

In this section, we describe Khovanov homology along the lines of [31, 3], and we tell the story so that the gradings and the structure of the differential emerge in a natural way. This approach to motivating the Khovanov homology uses elements of Khovanov's original approach, Viro's use of enhanced states for the bracket polynomial [48], and Bar-Natan's emphasis on tangle cobordisms [2, 3]. We use similar considerations in our paper [27].

Two key motivating ideas are involved in finding the Khovanov invariant. First of all, one would like to *categorify* a link polynomial such as $\langle K \rangle$. There are many meanings to the term categorify, but here the quest is to find a way to express the link polynomial as a *graded Euler characteristic* $\langle K \rangle = \chi_q \langle \mathcal{H}(K) \rangle$ for some homology theory associated with $\langle K \rangle$.

We will use the bracket polynomial and its enhanced states as described in the previous sections of this paper. To see how the Khovanov grading arises, consider the form of the expansion of this version of the bracket polynomial in enhanced states. We have the formula as a sum over enhanced states s :

$$\langle K \rangle = \sum_s (-1)^{i(s)} q^{j(s)}$$

where $i(s)$ is the number of B -type smoothings in s , $\lambda(s)$ is the number of loops in s labeled 1 minus the number of loops labeled X , and $j(s) = i(s) + \lambda(s)$. This can be rewritten in the following form:

$$\langle K \rangle = \sum_{i,j} (-1)^i q^j \dim(\mathcal{C}^{ij})$$

where we define \mathcal{C}^{ij} to be the linear span (over the complex numbers for the purpose of this paper, but over the integers or the integers modulo two for other contexts) of the set of enhanced states with $i(s) = i$ and $j(s) = j$. Then the number of such states is the dimension $\dim(\mathcal{C}^{ij})$.

We would like to have a bigraded complex composed of the \mathcal{C}^{ij} with a differential

$$\partial : \mathcal{C}^{ij} \longrightarrow \mathcal{C}^{i+1j}.$$

The differential should increase the *homological grading* i by 1 and preserve the *quantum grading* j . Then we could write

$$\langle K \rangle = \sum_j q^j \sum_i (-1)^i \dim(\mathcal{C}^{ij}) = \sum_j q^j \chi(\mathcal{C}^{\bullet j}),$$

where $\chi(\mathcal{C}^{\bullet j})$ is the Euler characteristic of the subcomplex $\mathcal{C}^{\bullet j}$ for a fixed value of j .

This formula would constitute a categorification of the bracket polynomial. Below, we shall see how *the original Khovanov differential ∂ is uniquely determined by the restriction that $j(\partial s) = j(s)$ for each enhanced state s* . Since j is preserved by the differential, these subcomplexes $\mathcal{C}^{\bullet j}$ have their own Euler characteristics and homology. We have

$$\chi(H(\mathcal{C}^{\bullet j})) = \chi(\mathcal{C}^{\bullet j})$$

where $H(\mathcal{C}^{\bullet j})$ denotes the homology of the complex $\mathcal{C}^{\bullet j}$. We can write

$$\langle K \rangle = \sum_j q^j \chi(H(\mathcal{C}^{\bullet j})).$$

The last formula expresses the bracket polynomial as a *graded Euler characteristic* of a homology theory associated with the enhanced states of the bracket state summation. This is the categorification of the bracket polynomial. Khovanov proves that this homology theory is an invariant of knots and links (via the Reidemeister moves of Figure 1), creating a new and stronger invariant than the original Jones polynomial.

We will construct the differential in this complex first for mod-2 coefficients. The differential is based on regarding two states as *adjacent* if one differs from the other by a single smoothing at some site. Thus if (s, τ) denotes a pair consisting in an enhanced state s and site τ of that state with τ of type A , then we consider all enhanced states s' obtained from s by smoothing at τ and relabeling only those loops that are affected by the resmoothing. Call this set of enhanced states $S'[s, \tau]$. Then we shall define the *partial differential* $\partial_\tau(s)$ as a sum over certain elements in $S'[s, \tau]$, and the differential by the formula

$$\partial(s) = \sum_\tau \partial_\tau(s)$$

with the sum over all type A sites τ in s . It then remains to see what are the possibilities for $\partial_\tau(s)$ so that $j(s)$ is preserved.

Note that if $s' \in S'[s, \tau]$, then $i(s') = i(s) + 1$. Thus

$$j(s') = i(s') + \lambda(s') = 1 + i(s) + \lambda(s').$$

From this we conclude that $j(s) = j(s')$ if and only if $\lambda(s') = \lambda(s) - 1$. Recall that

$$\lambda(s) = [s : +] - [s : -]$$

where $[s : +]$ is the number of loops in s labeled $+1$, $[s : -]$ is the number of loops labeled -1 (same as labeled with x) and $j(s) = i(s) + \lambda(s)$.

Proposition. The partial differentials $\partial_\tau(s)$ are uniquely determined by the condition that $j(s') = j(s)$ for all s' involved in the action of the partial differential on the enhanced state s . This unique form of the partial differential can be described by the following structures of multiplication and comultiplication on the algebra $V = k[x]/(x^2)$ where $k = \mathbb{Z}/2\mathbb{Z}$ for mod-2 coefficients, or $k = \mathbb{Z}$ for integral coefficients.

1. The element 1 is a multiplicative unit and $x^2 = 0$.
2. $\Delta(1) = 1 \otimes x + x \otimes 1$ and $\Delta(x) = x \otimes x$.

These rules describe the local relabeling process for loops in a state. Multiplication corresponds to the case where two loops merge to a single loop, while comultiplication corresponds to the case where one loop bifurcates into two loops.

Proof. Using the above description of the differential, suppose that there are two loops at τ that merge in the smoothing. If both loops are labeled 1 in s then the local contribution to $\lambda(s)$ is 2. Let s' denote a smoothing in $S[s, \tau]$. In order for the local λ contribution to become 1, we see that the merged loop must be labeled 1. Similarly if the two loops are labeled 1 and X , then the merged loop must be labeled X so that the local contribution for λ goes from 0 to -1 . Finally, if the two loops are labeled X and X , then there is no label available for a single loop that will give -3 , so we define ∂ to be zero in this case. We can summarize the result by saying that there is a multiplicative structure m such that $m(1, 1) = 1, m(1, x) = m(x, 1) = x, m(x, x) = 0$, and this multiplication describes the structure of the partial differential when two loops merge. Since this is the multiplicative structure of the algebra $V = k[x]/(x^2)$, we take this algebra as summarizing the differential.

Now consider the case where s has a single loop at the site τ . Smoothing produces two loops. If the single loop is labeled x , then we must label each of the two loops by x in order to make λ decrease by 1. If the single loop is labeled 1, then we can label the two loops by x and 1 in either order. In this second case we take the partial differential of s to be the sum of these two labeled states. This structure can be described by taking a coproduct structure with $\Delta(x) = x \otimes x$ and $\Delta(1) = 1 \otimes x + x \otimes 1$. We now have the algebra $V = k[x]/(x^2)$ with product $m : V \otimes V \rightarrow V$ and coproduct $\Delta : V \rightarrow V \otimes V$, describing the differential completely. This completes the proof. //

Partial differentials are defined on each enhanced state s and a site τ of type A in that state. We consider states obtained from the given state by smoothing the given site τ . The result of smoothing τ is to produce a new state s' with one more site of type B than s . Forming s' from s we either amalgamate two loops to a single loop at τ , or we divide a loop at τ into two distinct loops. In the case of amalgamation, the new state s acquires the label on the amalgamated circle that is the product of the labels on the two circles that are its ancestors in s . This case of the partial differential is described by the multiplication in the algebra. If one circle becomes two circles, then we apply the coproduct. Thus if the circle is labeled X , then the resultant two circles are each labeled X corresponding to $\Delta(x) = x \otimes x$. If the original circle is labeled 1 then we take

the partial boundary to be a sum of two enhanced states with labels 1 and x in one case, and labels x and 1 in the other case, on the respective circles. This corresponds to $\Delta(1) = 1 \otimes x + x \otimes 1$. Modulo two, the boundary of an enhanced state is the sum, over all sites of type A in the state, of the partial boundaries at these sites. It is not hard to verify directly that the square of the boundary mapping is zero (this is the identity of mixed partials!) and that it behaves as advertised, keeping $j(s)$ constant. There is more to say about the nature of this construction with respect to Frobenius algebras and tangle cobordisms. In Figures 4,5 and 6 we illustrate how the partial boundaries can be conceptualized in terms of surface cobordisms. Figure 4 shows how the partial boundary corresponds to a saddle point and illustrates the two cases of fusion and fission of circles. The equality of mixed partials corresponds to topological equivalence of the corresponding surface cobordisms, and to the relationships between Frobenius algebras [23] and the surface cobordism category. In particular, in Figure 6 we show how in a key case of two sites (labeled 1 and 2 in that Figure) the two orders of partial boundary are

$$\partial_2 \partial_1 = (1 \otimes m) \circ (\Delta \otimes 1)$$

and

$$\partial_1 \partial_2 = \Delta \circ m.$$

In the Frobenius algebra $V = k[x]/(x^2)$ we have the identity

$$(1 \otimes m) \circ (\Delta \otimes 1) = \Delta \circ m.$$

Thus the Frobenius algebra implies the identity of the mixed partials. Furthermore, in Figure 5 we see that this identity corresponds to the topological equivalence of cobordisms under an exchange of saddle points.

In Figures 7 and 8 we show another aspect of this algebra. As Figure 7 illustrates, we can consider cup (minimum) and cap (maximum) cobordisms that go between the empty set and a single circle. With the categorical arrow going down the page, the cap is a mapping from the base ring k to the module V and we denote this mapping by $\eta : k \rightarrow V$. It is the *unit* for the algebra V and is defined by $\eta(1) = 1_V$, taking 1 in k to 1_V in V . The cup is a mapping from V to k and is denoted by $\epsilon : V \rightarrow k$. This is the *counit*. As Figure 7 illustrates, we need a basic identity about the counit which reads

$$\sum \epsilon(a_1) a_2 = a$$

for any $a \in V$ where

$$\Delta(a) = \sum a_1 \otimes a_2.$$

The summation is over an appropriate set of elements in $v \otimes V$ as in our specific formulas for the algebra $k[x]/(x^2)$. Of course we also demand

$$\sum a_1 \epsilon(a_2) = a$$

for any $a \in V$. With these formulas about the counit and unit in place, we see that cobordisms will give equivalent algebra when one cancels a maximum or a minimum with a saddle point, again as shown in Figure 7.

Note that for our algebra $V = k[x]/(x^2)$, it follows from the counit identities of the last paragraph that

$$\epsilon(1) = 0$$

and

$$\epsilon(x) = 1.$$

In fact, Figure 8 shows a formula that holds in this special algebra. The formula reads

$$\epsilon(ab) = \epsilon(ax)\epsilon(b) + \epsilon(a)\epsilon(bx)$$

for any $a, b \in V$. As the rest of Figure 8 shows, this identity means that a single tube in any cobordism can be cut, replacing it by a cups and a caps in a linear combination of two terms. The tube-cutting relation is shown in its most useful form at the bottom of Figure 8. In Figure 8, the black dots are symbols standing for the special element x in the algebra.

It is important to note that we have a nonsingular pairing

$$\langle | \rangle : V \otimes V \longrightarrow k$$

defined by the equation

$$\langle a|b \rangle = \epsilon(ab).$$

One can define a Frobenius algebra by starting with the existence of a non-singular bilinear pairing. In fact, a finite dimensional associative algebra with unit defined over a unital commutative ring k is said to be a *Frobenius algebra* if it is equipped with a non-degenerate bilinear form

$$\langle | \rangle : V \otimes V \longrightarrow k$$

such that

$$\langle ab|c \rangle = \langle a|bc \rangle$$

for all a, b, c in the algebra. The other mappings and the interpretation in terms of cobordisms can all be constructed from this definition. See [23].

Remark on Grading and Invariance. In Section 2 we showed how the bracket, using the variable q , behaves under Reidemeister moves. These formulas correspond to how the invariance of the homology works in relation to the moves. We have that

$$J_K = (-1)^{n_-} q^{n_+ - 2n_-} \langle K \rangle,$$

where n_- denotes the number of negative crossings in K and n_+ denotes the number of positive crossings in K . $J(K)$ is invariant under all three Reidemeister moves. The corresponding formulas for Khonavov homology are as follows

$$J_K = (-1)^{n_-} q^{n_+ - 2n_-} \langle K \rangle =$$




Figure 4: **SaddlePoints and State Smoothings**

$$\begin{aligned}
& (-1)^{n-} q^{n+-2n-} \sum_{i,j} (-1)^i a^j \dim(H^{i,j}(K)) = \\
& \sum_{i,j} (-1)^{i+n+} q^{j+n+-2n-1} \dim(H^{i,j}(K)) = \\
& \sum_{i,j} (-1)^i q^j \dim(H^{i-n-,j-n++2n-}(K)).
\end{aligned}$$

It is often more convenient to define the *Poincaré polynomial* for Khovanov homology via

$$P_K(t, q) = \sum_{i,j} t^i q^j \dim(H^{i-n-,j-n++2n-}(K)).$$

The Poincaré polynomial is a two-variable polynomial invariant of knots and links, generalizing the Jones polynomial. Each coefficient

$$\dim(H^{i-n-,j-n++2n-}(K))$$

is an invariant of the knot, invariant under all three Reidemeister moves. In fact, the homology groups

$$H^{i-n-,j-n++2n-}(K)$$

are knot invariants. The grading compensations show how the grading of the homology can change from diagram to diagram for diagrams that represent the same knot.

Remark on Integral Differentials. Choose an ordering for the crossings in the link diagram K and denote them by $1, 2, \dots, n$. Let s be any enhanced state of K and let $\partial_i(s)$ denote the chain obtained from s by applying a partial boundary at the i -th site of s . If the i -th site is a smoothing of type A^{-1} , then $\partial_i(s) = 0$. If the i -th site is a smoothing of type A , then $\partial_i(s)$ is given by the rules discussed above (with the same signs). The compatibility conditions that we have discussed show that partials commute in the sense that $\partial_i(\partial_j(s)) = \partial_j(\partial_i(s))$ for all i and j . One then defines





Figure 5: **Surface Cobordisms**



$$\partial_2 \partial_1 = (1 \otimes m) (\Delta \otimes 1)$$

$$\partial_1 \partial_2 = (\Delta) (m)$$

$$\partial_1 \partial_2 = \partial_2 \partial_1$$

Figure 6: **Local Boundaries Commute**




Figure 7: Unit and Counit as Cobordisms




Figure 8: The Tube Cutting Relation

signed boundary formulas in the usual way of algebraic topology. One way to think of this regards the complex as the analogue of a complex in de Rham cohomology. Let $\{dx_1, dx_2, \dots, dx_n\}$ be a formal basis for a Grassmann algebra so that $dx_i \wedge dx_j = -dx_j \wedge dx_i$. Starting with enhanced states s in $C^0(K)$ (that is, states with all A -type smoothings) define formally, $d_i(s) = \partial_i(s)dx_i$ and regard $d_i(s)$ as identical with $\partial_i(s)$ as we have previously regarded it in $C^1(K)$. In general, given an enhanced state s in $C^k(K)$ with B -smoothings at locations $i_1 < i_2 < \dots < i_k$, we represent this chain as $s dx_{i_1} \wedge \dots \wedge dx_{i_k}$ and define

$$\partial(s dx_{i_1} \wedge \dots \wedge dx_{i_k}) = \sum_{j=1}^n \partial_j(s) dx_j \wedge dx_{i_1} \wedge \dots \wedge dx_{i_k},$$

just as in a de Rham complex. The Grassmann algebra automatically computes the correct signs in the chain complex, and this boundary formula gives the original boundary formula when we take coefficients modulo two. Note, that in this formalism, partial differentials ∂_i of enhanced states with a B -smoothing at the site i are zero due to the fact that $dx_i \wedge dx_i = 0$ in the Grassmann algebra. There is more to discuss about the use of Grassmann algebra in this context. For example, this approach clarifies parts of the construction in [28].

It of interest to examine this analogy between the Khovanov (co)homology and de Rham cohomology. In that analogy the enhanced states correspond to the differentiable functions on a manifold. The Khovanov complex $C^k(K)$ is generated by elements of the form $s dx_{i_1} \wedge \dots \wedge dx_{i_k}$ where the enhanced state s has B -smoothings at exactly the sites i_1, \dots, i_k . If we were to follow the analogy with de Rham cohomology literally, we would define a new complex $DR(K)$ where $DR^k(K)$ is generated by elements $s dx_{i_1} \wedge \dots \wedge dx_{i_k}$ where s is *any* enhanced state of the link K . The partial boundaries are defined in the same way as before and the global boundary formula is just as we have written it above. This gives a *new* chain complex associated with the link K . Whether its homology contains new topological information about the link K will be the subject of a subsequent paper.

In the case of de Rham cohomology, we can also look for compatible unitary transformations. Let M be a differentiable manifold and $\mathcal{C}(M)$ denote the DeRham complex of M over the complex numbers. Then for a differential form of the type $f(x)\omega$ in local coordinates x_1, \dots, x_n and ω a wedge product of a subset of $dx_1 \dots dx_n$, we have

$$d(f\omega) = \sum_{i=1}^n (\partial f / \partial x_i) dx_i \wedge \omega.$$

Here d is the differential for the DeRham complex. Then $\mathcal{C}(M)$ has as basis the set of $|f(x)\omega\rangle$ where $\omega = dx_{i_1} \wedge \dots \wedge dx_{i_k}$ with $i_1 < \dots < i_k$. We could achieve $Ud + dU = 0$ if U is a very simple unitary operator (e.g. multiplication by phases that do not depend on the coordinates x_i) but in general it will be an interesting problem to determine all unitary operators U with this property.

A further remark on de Rham cohomology. There is another relation with the de Rham complex: In [41] it was observed that Khovanov homology is related to Hochschild homology and Hochschild homology is thought to be an algebraic version of de Rham chain complex (cyclic cohomology corresponds to de Rham cohomology), compare [45].

5 The Cube Category and the Tangle Cobordism Structure of Khovanov Homology

We can now connect the constructions of the last section with the homology construction via the cube category. Here it will be convenient to think of the state category $\mathcal{S}(K)$ as a cube category with extra structure. Thus we will denote the bracket states by sequences of A 's and B 's as in Figures 2 and 3. And we shall regard the maps such as $d_2 : \langle AABA \rangle \longrightarrow \langle ABBA \rangle$ as corresponding to re-smoothings of bracket states that either join or separate state loops. We take $V = k[x]/(x^2)$ with the coproduct structure as given in the previous section. The maps from $m : V \otimes V \longrightarrow V$ and $\Delta : V \longrightarrow V \otimes V$ allow us to define the images of the resmoothing maps d_i under a functor $\mathcal{F} : \mathcal{S}(K) \longrightarrow \mathcal{M}$ where \mathcal{M} is the category generated by V by taking tensor powers of V and direct sums of these tensor powers. It then follows that the homology we have described in the previous section is exactly the homology associated with this functor.

The material in the previous section also suggests a modification of the state category $\mathcal{S}(K)$. Instead of taking the maps in this category to be simply the abstract arrows generated by elementary re-smoothings of states from A to B , we can regard each such smoothing as a surface cobordism from the set of circles comprising the domain state to the set of circles comprising the codomain state. With this, in mind, two such cobordisms represent equivalent morphisms whenever the corresponding surfaces are homeomorphic relative to their boundaries. Call this category $\mathcal{CobS}(K)$. We then easily generalize the observations of the previous section, particularly Figures 4, 5 and 6, to see that we have the desired commuting relations $d_i d_j = d_j d_i$ (for $i \neq j$) in $\mathcal{CobS}(K)$ so that any functor from $\mathcal{CobS}(K)$ to a module category will have a well-defined chain complex and associated homology. This applies, in particular to the functor we have constructed, using the Frobenius algebra $V = k[x]/(x^2)$.

In [3] BarNatan takes the approach using surface cobordisms a step further by making a categorical analog of the chain complex. For this purpose we let $\mathcal{CobS}(K)$ become an additive category. Maps between specific objects X and Y added formally and the set $\mathit{Maps}(X, Y)$ is a module over the integers. More generally, let \mathcal{C} be an additive category. In order to create the analog of a chain complex, let $\mathit{Mat}(\mathcal{C})$ denote the *Matrix Category of \mathcal{C}* whose objects are n -tuples (vectors) of objects of \mathcal{C} (n can be any natural number) and whose morphisms are in the form of a matrix $m = (m_{ij})$ of morphisms in \mathcal{C} where we write

$$m : O \longrightarrow O'$$

and

$$m_{ij} : O_i \longrightarrow O'_j$$

for

$$\begin{aligned} O &= (O_1, \dots, O_n), \\ O' &= (O'_1, \dots, O'_m). \end{aligned}$$

Here O_i and O'_j are objects in \mathcal{C} while O and O' are objects in $\mathcal{M}at(\mathcal{C})$. Composition of morphisms in $\mathcal{M}at(\mathcal{C})$ follows the pattern of matrix multiplication. If

$$n : O' \longrightarrow O''$$

then

$$n \circ m : O \longrightarrow O''$$

and

$$(n \circ m)_{i,j} = \sum_k n_{i,k} \circ m_{k,j}$$

where the compositions in the summation occur in the category \mathcal{C} .

We then define the *category of complexes over \mathcal{C}* , denoted $\mathcal{K}om(\mathcal{M}at(\mathcal{C}))$ to consist of sequences of objects of $\mathcal{M}at(\mathcal{C})$ and maps between them so that consecutively composed maps are equal to zero.

$$\dots \longrightarrow O^k \longrightarrow O^{k+1} \longrightarrow O^{k+2} \longrightarrow \dots$$

Here we let $\partial_k : O^k \longrightarrow O^{k+1}$ denote the differential in the complex and we assume that $\partial_{k+1}\partial_k = 0$. A morphism between complexes O^* and O'^* consists in a family of maps $f_k : O^k \longrightarrow O'^k$ such that $\partial'_k f_k = f_{k+1} \partial_k$. Such morphisms will be called *chain maps*.

At this abstract level, we cannot calculate homology since kernels and cokernels are not available, but we can define the *homotopy type* of a complex in $\mathcal{K}om(\mathcal{M}at(\mathcal{C}))$. We say that two chain maps $f : O \longrightarrow O'$ and $g : O \longrightarrow O'$ are *homotopic* if there is a sequence of mappings $H_k : O^k \longrightarrow O^{k-1}$ such that

$$f - g = H\partial + \partial H.$$

Specifically, this means that

$$f_k - g_k = H_{k+1}\partial_k + \partial_{k+1}H_k.$$

Note that if $\phi = H\partial + \partial H$, then

$$\partial\phi = \partial H\partial = \phi\partial.$$

Thus any such ϕ is a chain map. We call two complexes O and O' *homotopy equivalent* if there are chain maps $F : O \longrightarrow O'$ and $G : O' \longrightarrow O$ such that both FG and GF are homotopic to the identity map of O and O' respectively. The homotopy type of a complex is an abstract substitute for the homology since, in an abelian category (where one can compute homology) homology is an invariant of homotopy type.

We are now in a position to work with the category $\mathcal{K}om(\mathcal{M}at(\mathcal{C}obS(K)))$ where K is a link diagram. The question is, what extra equivalence relation on the category $\mathcal{C}obS(K)$ will ensure that the homotopy types in $\mathcal{K}om(\mathcal{M}at(\mathcal{C}obS(K)))$ will be invariant under Reidemeister moves on the diagram K .

BarNatan [3] gives an elegant answer to this question. His answer is illustrated in Figure 9 where we show the *4Tu Relation*, the *Sphere Relation* and the *Torus Relation*. The key relation is the the *4Tu* relation, which serves a number of purposes, including being a basic homotopy in the category $\mathcal{K}om(\mathcal{M}at(\mathcal{C}obS(K)))$. Note that the *4Tu* relation can be described as follows: There are four local bits of surface, call them S_1, S_2, S_3, S_4 . Let $C_{i,j}$ denote this configuration with a tube connecting S_i and S_j . Then in the cobordism category we take the identity

$$C_{1,2} + C_{3,4} = C_{1,3} + C_{2,4}.$$

It is a good exercise for the reader to show that the *4Tu* relation follows from the tube cutting relation of Figure 8. The Sphere and Torus relations assert that the 2-sphere has value 0 and that the torus has value 2, just as we have seen by using the Frobenius algebra in Figure 7.

To illustrate how things work once we factor by these relations, we show in Figures 10 and 11 how one sees the homotopy equivalence of the complexes for a diagram before and after the second Reidemeister move. In Figure 10 we show the complexes and indicate chain maps F and G between them and homotopies in the complex for the diagram before it is simplified by the Reidemeister move. In Figure 11 we show these maps and their compositions in the form of a four-term identity. The reader should first just look at the identity of involving these four cobordisms and recognize that it is exactly the *4Tu* relation. On the other hand, each of the terms in the relation is factored into mappings involving F_1, G_1 and the homotopies H_1 and H_2 and the boundary mappings in the complex. Study of Figure 11 will convince the reader that the complexes before and after the second Reidemeister move are homotopy equivalent. A number of details are left to the reader. For example, note that in Figure 10 we have indicated the categorical chain complexes Z and W by showing only how they differ locally near the change corresponding to a Reidemeister two move. We give, via Figures 10 and 11, chain maps $F : W \rightarrow Z$ and $G : Z \rightarrow W$. These maps consist in a particular cobordism on one part of the complex and an identity map on the other part of the complex. We have specifically labeled parts of these mappings by F_1 and by G_1 . Using the implicit definitions of F_1 and G_1 given in Figure 11, the reader will easily see that $G_1 F_1 = 0$ since this composition includes a 2-sphere. From this it follows that GF is the identity mapping on the complex W . We also leave to the reader to check that the mappings F and G commute with the boundary mappings so that they are mappings of complexes. The part of the homotopy indicated shows that FG is homotopic to the identity (up to sign) and so shows that the complexes Z and W are homotopy equivalent. One needs the value of the torus equal to 2 for homotopy invariance under the first Reidemeister move. This is a fragment of the full derivation of homotopy equivalences corresponding to all three Reidemeister moves that is given in the paper [3].




Figure 9: **The 4Tu Relation, Sphere and Torus Relations**




Figure 10: **Complexes for Second Reidemeister Move**




Figure 11: Homotopy for Second Reidemeister Move

6 Other Frobenius Algebras and Rasmussen's Theorem

Lee [25] makes another homological invariant of knots and links by using a different Frobenius algebra. She takes the algebra $\mathcal{A} = k[x]/(x^2 - 1)$ with

$$x^2 = 1,$$

$$\Delta(1) = 1 \otimes x + x \otimes 1,$$

$$\Delta(x) = x \otimes x + 1 \otimes 1,$$

$$\epsilon(x) = 1,$$

$$\epsilon(1) = 0.$$

This gives a link homology theory that is distinct from Khovanov homology. In this theory, the quantum grading j is not preserved, but we do have that

$$j(\partial(\alpha)) \geq j(\alpha)$$

for each chain α in the complex. This means that *one can use j to filter the chain complex for the Lee homology*. The result is a spectral sequence that starts from Khovanov homology and converges to Lee homology.

Lee homology is simple. One has that the dimension of the Lee homology is equal to $2^{\text{comp}(L)}$ where $\text{comp}(L)$ denotes the number of components of the link L . Up to homotopy, Lee's homology has a vanishing differential, and the complex behaves well under link concordance. In his paper [4] Dror BarNatan remarks "In a beautiful article Eun Soo Lee introduced a second differential on the Khovanov complex of a knot (or link) and showed that the resulting (double) complex has non-interesting homology. This is a very interesting result." Rasmussen [44] uses Lee's result to define invariants of links that give lower bounds for the four-ball genus, and determine it for torus knots. This gives an (elementary) proof of a conjecture of Milnor that had been previously shown using gauge theory by Kronheimer and Mrowka [24].

Rasmussen's result uses the Lee spectral sequence. We have the quantum (j) grading for a diagram K and the fact that for Lee's algebra $j(\partial(s)) \geq j(s)$. Rasmussen uses a normalized version of this grading denoted by $g(s)$. Then one makes a filtration $F^k C^*(K) = \{v \in C^*(K) | g(v) \geq k\}$ and given $\alpha \in \text{Lee}^*(K)$ define

$$S(\alpha) := \max\{g(v) | [v] = \alpha\}$$

$$s_{\min}(K) := \min\{S(\alpha) | \alpha \in \text{Lee}^*(K), \alpha \neq 0\}$$

$$s_{\max}(K) := \max\{S(\alpha) | \alpha \in \text{Lee}^*(K), \alpha \neq 0\}$$

and

$$s(K) := (1/2)(s_{\min}(K) + s_{\max}(K)).$$

This last average of s_{\min} and s_{\max} is the Rasmussen invariant.

We now enter the following sequence of facts:

1. $s(K) \in \mathbb{Z}$.
2. $s(K)$ is additive under connected sum.
3. If K^* denotes the mirror image of the diagram K , then

$$s(K^*) = -s(K).$$

4. If K is a positive knot diagram (all positive crossings), then

$$s(K) = -r + n + 1$$

where r denotes the number of loops in the canonical oriented smoothing (this is the same as the number of Seifert circuits in the diagram K) and n denotes the number of crossings in K .

5. For a torus knot $K_{a,b}$ of type (a, b) , $s(K_{a,b}) = (a - 1)(b - 1)$.
6. $|s(K)| \leq 2g^*(K)$ where $g^*(K)$ is the least genus spanning surface for K in the four ball.
7. $g^*(K_{a,b}) = (a - 1)(b - 1)/2$. This is Milnor's conjecture.

This completes a very skeletal sketch of the construction and use of Rasmussen's invariant.

7 The Simplicial Structure of Khovanov Homology

Let \mathcal{S} denote the set of (standard) bracket states for a link diagram K . One way to describe the Khovanov complex is to associate to each state loop λ a module V isomorphic to the algebra $k[x]/(x^2)$ with coproduct as we have described in the previous sections. The generators 1 and x of this algebra can then be regarded as the two possible enhancements of the loop λ . In the same vein we associate to a state S the tensor product of copies of V , one copy for each loop in the state. The local boundaries are defined exactly as before, and the Khovanov complex is the direct sum of the modules associated with the states of the link diagram. We will use this point of view in the present section, and we shall describe Khovanov homology in terms of the n -cube category and an associated simplicial object. The purpose of this section is to move towards, albeit in an abstract manner, a description of Khovanov homology as the homology of a topological space whose homotopy type is an invariant of the knot of the underlying knot or link. We do not accomplish this aim, but the constructions given herein may move toward that goal. An intermediate possibility would be to replace the Khovanov homology by an abstract space or simplicial object whose generalized homotopy type was an invariant of the knot or link.

Let $\mathcal{D}^n = \{A, B\}^n$ be the n -cube category whose objects are the n -sequences from the set $\{A, B\}$ and whose morphisms are arrows from sequences with greater numbers of A 's to sequences with fewer numbers of A 's. Thus \mathcal{D}^n is equivalent to the poset category of subsets of $\{1, 2, \dots, n\}$. Let \mathcal{M} be a pointed category with finite sums, and let $\mathcal{F} : \mathcal{D}^n \rightarrow \mathcal{M}$ be a functor. In our case \mathcal{M} is a category of modules (as described above) and \mathcal{F} carries n -sequences to certain tensor powers corresponding to the standard bracket states of a knot or link K . We map sequences to states by choosing to label the crossings of the diagram K from the set $\{1, 2, \dots, n\}$, and letting the functor take abstract A 's and B 's in the cube category to smoothings at those crossings of type A or type B . Thus each sequence in the cube category is associated with a unique state of K when K has n crossings. Nevertheless, we shall describe the construction more generally.

For the functor \mathcal{F} we first construct a semisimplicial object $\mathcal{C}(F)$ over \mathcal{M} , where a semisimplicial object is a simplicial object without degeneracies. This means that it has partial boundaries analogous to the partial boundaries that we have discussed before but none of the degeneracy maps that are common to simplicial theory (see [46] Chapter 1). For $k \geq 0$ we set

$$\mathcal{C}(F)_k = \bigoplus_{v \in \mathcal{D}_k^n} \mathcal{F}(v)$$

where \mathcal{D}_k^n denotes those sequences in the cube category with k A 's. Note that we are indexing dually to the upper indexing in the Khovanov homology sections of this paper where we counted the number of B 's in the states.

We introduce face operators (partial boundaries in our previous terminology)

$$d_i : \mathcal{C}(F)_k \rightarrow \mathcal{C}(F)_{k-1}$$

for $0 \leq i \leq k$ with $k \geq 1$ as follows: d_i is trivial for $i = 0$ and otherwise d_i acts on $\mathcal{F}(v)$ by the map $\mathcal{F}(v) \rightarrow \mathcal{F}(v')$ where v' is the sequence resulting from replacing the i -th A by B . The operators d_i satisfy the usual face relations of simplicial theory:

$$d_i d_j = d_{j-1} d_i$$

for $i < j$.

We now expand $\mathcal{C}(F)$ to a simplicial object $\mathcal{S}(F)$ over \mathcal{M} by applying freely degeneracies to the $\mathcal{F}(v)$'s. Thus

$$\mathcal{S}(F)_m = \bigoplus_{v \in \mathcal{D}_k^n, k+t=m} s_{i_1} \cdots s_{i_t} \mathcal{F}(v)$$

where $m > i_1 > \cdots > i_t \geq 0$ and these degeneracy operators are applied freely modulo the usual (axiomatic) relations among themselves and with the face operators. Then $\mathcal{S}(F)$ has degeneracies via formal application of degeneracy operators to these forms, and has face operators extending those of $\mathcal{C}(F)$. It is at this point we should remark that in our knot theoretic construction there is only at this point an opportunity for formal extension of degeneracy operators above the number of crossings in the given knot or link diagram since to make specific degeneracies would involve the creation of new diagrammatic sites. There may be a natural construction of this sort and if there is, such a construction could lead to a significant homotopy theory for Khovanov homology.

When the functor $\mathcal{F} : \mathcal{D}^n \longrightarrow \mathcal{M}$ goes to an abelian category \mathcal{M} , as in our knot theoretic case, we can recover the homology groups via

$$H_*NS(\mathcal{F}) \cong H_*\mathcal{C}(\mathcal{F})$$

where $NS(\mathcal{F})$ is the normalized chain complex of $\mathcal{S}(\mathcal{F})$. This completes the abstract simplicial description of this homology.

8 Quantum Comments

States of a quantum system are represented by unit vectors in a Hilbert space. Quantum processes are unitary transformations applied to these state vectors. In an appropriate basis for the Hilbert space, each basis vector represents a possible measurement. If $|\psi\rangle$ is a unit vector, then, upon measurement, one of the basis vectors will appear with probability, the absolute square of its coefficient in $|\psi\rangle$. One can, in principle, find the trace of a given unitary transformation by instantiating it in a certain quantum system and making repeated measurements on that system. Such a scheme, in the abstract, is called a quantum algorithm, and in the concrete is called a quantum computer. One well-known quantum algorithm for determining the trace of a unitary matrix is called the ‘‘Hadamard Test’’ [47].

In [19] we consider the Jones polynomial and Khovanov homology in a quantum context. In this section we give a sketch of these ideas. Recall from Section 2 that we have the following formula for the Jones polynomial.

$$J_K = (-1)^{n_-} q^{n_+ - 2n_-} \langle K \rangle.$$

Using the enhanced states formulation of Section 2, we form a Hilbert space $\mathcal{H}(K)$ with orthonormal basis the set of enhanced states of K . For the Hilbert space we denote a basis element by $|s\rangle$ where s is an enhanced state of the diagram K . Now using q as in Section 2, let q be any point on the unit circle in the complex plane. Define $U_K : \mathcal{H}(K) \longrightarrow \mathcal{H}(K)$ by the formula

$$U_K |s\rangle = (-1)^{i(s)+n_-} q^{j(s)+n_+ - 2n_-} |s\rangle.$$

Then U_K defines a unitary transformation of the Hilbert space and we have that

$$J_K = \text{Trace}(U_K).$$

The Hadamard Test applied to this unitary transformation gives a quantum algorithm for the Jones polynomial. This is not the most efficient quantum algorithm for the Jones polynomial. Unitary braid group representations can do better [20, 21, 1]. But this algorithm has the conceptual advantage of being directly related to Khovanov homology. In particular, let $C^{i,j}$ be the subspace of $\mathcal{H}(K)$ with basis the set of enhanced states $|s\rangle$ with $i(s) = i$ and $j(s) = j$. Then $\mathcal{H}(K)$ is the direct sum of these subspaces and we see that $\mathcal{H}(K)$ is identical with the Khovanov complex for

K with coefficients in the complex numbers. Furthermore, letting $\partial : \mathcal{H}(K) \rightarrow \mathcal{H}(K)$ be the boundary mapping that we have defined for the Khovanov complex, we have

$$\partial \circ U_K + U_K \circ \partial = 0.$$

Thus U_K induces a mapping on the Khovanov homology of K . As a linear space, the Khovanov homology of K ,

$$\text{Homology}(\mathcal{H}(K)) = \text{Kernel}(\partial) / \text{Image}(\partial)$$

is also a Hilbert space on which U_K acts and for which the trace yields the Jones polynomial.

If we are given more information about the Khovanov homology as a space, for example if we are given a basis for $H^{i-n_-, j-n_+ + 2n_-}(K)$ for each i and j , then we can extend U to act on $H^{i-n_-, j-n_+ + 2n_-}(K)$ as an eigenspace with eigenvalue $t^i q^j$ where q and t are chosen unit complex numbers. Then we have an extended U'_K with

$$U'_K |\alpha\rangle = t^i q^j |\alpha\rangle$$

for each $\alpha \in H^{i-n_-, j-n_+ + 2n_-}(K)$. With this extension we have that the trace of U'_K recovers a specialization of the Poincaré polynomial (Section 4) for the Khovanov homology.

$$\text{Trace}(U'_K) = \sum_{i,j} t^i q^j \dim(H^{i-n_-, j-n_+ + 2n_-}(K)) = P_K(t, q).$$

Thus, in principle, we formulate a quantum algorithm for specializations of the Poincaré polynomial for Khovanov homology.

Placing Khovanov homology in an appropriate quantum mechanical, quantum information theoretic, or quantum field theory context is a fundamental question that has been considered by a number of people, including Sergei Gukov [10, 11] and Edward Witten [52, 53, 54]. The constructions discussed here are elementary in nature but we would like to know how they interface with other points of view. In particular, if one thinks of the states in the state expansion of the bracket polynomial as analogs of the states of a physical system such as the Potts model in statistical mechanics, then the loop configuration of a given state corresponds to a decomposition of the underlying graph of the statistical mechanics model into regions of constant spin (where spin designates the local variable in the model). Working with a boundary operator, as we did with the Khovanov chain complex, means taking into account adjacency relations among these types of physical states.

9 Discussion

The subject of Khovanov homology is part of the larger subject of categorification in general and other link homologies in particular. The term *categorification* was coined by Crane and Frenkel in their paper [9] speculating on the possibility for invariants of four-manifolds via a categorical

generalization of Hopf algebras where all structures are moved up one categorical level. Just such a shift is seen in the Khovanov homology where loops that were once scalars become modules and the original Jones polynomial is seen as a graded Euler characteristic of a homology theory. There is now a complex literature on categorifications of quantum groups (aka Hopf algebras) and relationships of this new form of representation theory with the construction of link homology. For this we refer the reader to the following references [7, 8, 32, 33, 34, 43, 49, 50, 51]. It is possible that the vision of Crane and Frenkel for the construction of invariants of four dimensional manifolds will come true.

Other link homology theories are worth mentioning. In [35, 36, 37] Khovnaov and Rozansky construct a link homology theory for specializations of the Homflypt polynomial. Their theory extends integrally to a Khovanov homology theory for virtual knots, but no calculations are known at this writing. Khovanov homology does extend integrally to virtual knot theory as shown by Manturov in [28]. The relationship of the Manturov construction to that of Khovanov and Rozansky is not known at this time. In [27] Dye, Kauffman and Maturov show how to modify mod-2 Khovanov homology to categorify the arrow polynomial for virtual knots. This leads to many new calculations and examples [29, 30]. In [38, 39] Ozsvath, Szabo, Sarkar and Thurston construct combinatorial link homology based on Floer homology that categorifies the Alexander polynomial. Their techniques are quite different from those explained here for Khovanov homology. The combinatorial definition should be compared with that of Khovanov homology, but it has a flavor that is different, probably due to the fact that it categorifies a determinant that calculates the Alexander polynomial. This Knot Floer Homology theory is very powerful and can detect the three-dimensional genus of a knot (the least genus of an orientable spanning surface for the knot in three dimensional space). Caprau in [5] has a useful version of the tangle cobordism approach to Khovanov homology and Clark, Morisson and Walker [6] have an oriented tangle cobordism theory that is used to sort out the functoriality of Khovanov homology for knot cobordisms. There is another significant variant of Khovanov homology termed *odd Khovanov homology* [40]. Attempts to find other global interpretations of Khovanov homology have led to very significant lines of research [7, 8, 43], and attempts to find general constructions for link homology corresponding to the quantum link invariants coming from quantum groups have led to research such as that of Webster [50, 51] where we now have theories for such constructions that use the categorifications of quantum groups for classical Lie algebras..

There have been three applications of Khovanov homology that are particularly worth mentioning. One, we have discussed in Section 6, is Rasmussen's use of Khovanov homology [44] to determine the slice genus of torus knots without using gauge theory. Another is the proof by Kronheimer and Mrowka [24] that Khovanov homology detects the unknot. The work of Kronheimer and Mrowka interrelates Khovanov homology with their theory of knot instanton homology and allows them to apply their gauge theoretic results to obtain this striking result. A proof that Khovanov homology detects the unknot by purely combinatorial topological means is unknown at this writing. By the same token, it is still unknown whether the Jones polynomial detects classical knots. Finally, we mention the work of Shumakovitch [42] where, by calculating Khovanov homology, he shows many examples of knots that are topologically slice but are not slice in the

differentiable category. Here Khovanov homology circumvents a previous use of gauge theory.

References

- [1] D. Aharonov, V. Jones, Z. Landau, A polynomial quantum algorithm for approximating the Jones polynomial, [quant-ph/0511096](#).
- [2] D. Bar-Natan, On Khovanov's categorification of the Jones polynomial, *Algebra Geometry Topology* Vol 2. (2002) 337-370, [math.QA/0201043](#).
- [3] D. Bar-Natan (2005), Khovanov's homology for tangles and cobordisms, *Geometry and Topology*, **9-33**, pp. 1465-1499. [arXiv:math.GT/0410495](#)
- [4] D. Bar-Natan and S. Morrison, The Karoubi envelope and Lee's degeneration of Khovanov homology. *Algebr. Geom. Topol.* 6 (2006), 14591469. [math.GT/0606542](#).
- [5] C. Caprau, The universal $\mathfrak{sl}(2)$ cohomology via webs and foams, *Topology and its Applications*, Volume 156 (2009), 1684-1702. [math.GT.arXiv:0802.2848](#)
- [6] D. Clark, S. Morrison, K. Walker, Fixing the functoriality of Khovanov homology. *Geom. Topol.* 13 (2009), no. 3, 14991582.
- [7] S. Cautis, J. Kamnitzer, Knot homology via derived categories of coherent sheaves. I. The $\mathfrak{sl}(2)$ -case. *Duke Math. J.* 142 (2008), no. 3, 511588.
- [8] S. Cautis, J. Kamnitzer, Knot homology via derived categories of coherent sheaves. II. slm case. *Invent. Math.* 174 (2008), no. 1, 165232.
- [9] L. Crane, I. Frenkel, Four-dimensional topological quantum field theory, Hopf categories, and the canonical bases. *Topology and physics. J. Math. Phys.* 35 (1994), no. 10, 51365154.
- [10] S. Gukov, Gauge theory and knot homologies. *Fortschr. Phys.* 55 (2007), no. 5-7, 473490.
- [11] S. Gukov, Surface operators and knot homologies, [arXiv:0706.2369](#).
- [12] V.F.R. Jones, A polynomial invariant for links via von Neumann algebras, *Bull. Amer. Math. Soc.*, **129** (1985), 103–112.
- [13] V.F.R. Jones. Hecke algebra representations of braid groups and link polynomials, *Ann. of Math.* **126**, (1987), pp. 335-338.
- [14] V.F.R. Jones, On knot invariants related to some statistical mechanics models, *Pacific J. Math.*, **137**, no. 2 (1989), pp. 311-334.

- [15] L.H. Kauffman, State models and the Jones polynomial, *Topology*, **26** (1987), 395–407.
bibitemKaAM L. H. Kauffman, New invariants in knot theory, Amer. Math. Monthly (1987).
- [16] L.H. Kauffman, Statistical mechanics and the Jones polynomial, *AMS Contemp. Math. Series*, **78** (1989), 263–297.
- [17] L.H. Kauffman, *Knots and Physics*, World Scientific Publishers (1991), Second Edition (1993), Third Edition (2002).
- [18] L.H. Kauffman, Topological quantum information, Khovanov homology and the Jones polynomial. Topology of algebraic varieties and singularities, 245264, *Contemp. Math.*, 538, Amer. Math. Soc., Providence, RI, 2011 [arXiv:1001.0354](https://arxiv.org/abs/1001.0354)
- [19] L. H. Kauffman. A quantum model for the Jones polynomial, Khovanov Homology and Generalized Simplicial Homology, *AMS Contemp Math*, **536**, Cross Disciplinary Advances in Quantum Computing, ed. by Mahdavi et al., pp. 75 - 94.
- [20] L.H. Kauffman, Quantum computing and the Jones polynomial, in *Quantum Computation and Information*, S. Lomonaco, Jr. (ed.), AMS CONM/305, 2002, pp. 101–137. [math.QA/0105255](https://arxiv.org/abs/math.QA/0105255)
- [21] L. H. Kauffman and S. Lomonaco Jr., A Three-stranded quantum algorithm for the Jones polynomial, in *Quantum Information and Quantum Computation V*, Proceedings of Spie, April 2007, edited by E.J. Donkor, A.R. Pirich and H.E. Brandt, pp. 65730T1-17, Intl Soc. Opt. Eng.
- [22] M. Khovanov, A categorification of the Jones polynomial, *Duke J. Math.* 101(200), no. 3, 359-426, [mathQA/9908171](https://arxiv.org/abs/mathQA/9908171).
- [23] J. Kock, “Frobenius Algebras and 2D Topological Quantum Field Theories”, London Mathematical Society, Student Texts No. 59 (2004).
- [24] P. B. Kronheimer, T. S. Mrowka, Khovanov homology is an unknot-detector. [math.GT.arXiv:0005046](https://arxiv.org/abs/math.GT/0005046).
- [25] E. S. Lee, An endomorphism of the Khovanov invariant, *Adv. Math.* 197 (2), 554-586 (2005). [math.GT/0210213](https://arxiv.org/abs/math.GT/0210213).
- [26] S. J. Lomonaco Jr. and L. H. Kauffman, Quantum Knots and Mosaics, *Journal of Quantum Information Processing*, Vol. 7, Nos. 2-3, (2008), pp. 85 - 115. arxiv.org/abs/0805.0339
- [27] H.A. Dye, L.H. Kauffman, V.O. Manturov, On two categorifications of the arrow polynomial for virtual knots, [arXiv:0906.3408](https://arxiv.org/abs/0906.3408).

- [28] V.O. Manturov, Khovanov homology for virtual links with arbitrary coefficients, *math.GT/0601152*. (Russian) *Izv. Ross. Akad. Nauk Ser. Mat.* 71 (2007), no. 5, 111–148. *J. Knot Theory Ramifications* 16 (2007), no. 3, 345377.
- [29] A. Kaestner Ph.D. Thesis, University of Illinois at Chicago (2011).
- [30] A. Kasetner and L. H. Kauffman, On Categorifications of the Arrow Polynomial (to appear).
- [31] M. Khovanov,(1997), A categorification of the Jones polynomial, *Duke Math. J.***101** (3), pp.359-426.
- [32] M. Khovanov, A. Lauda, A categorification of quantum $sl(n)$. *Quantum Topol.* 1 (2010), no. 1, 192.
- [33] M. Khovanov, A. Lauda, A diagrammatic approach to categorification of quantum groups II.*Trans. Amer. Math. Soc.* 363 (2011), no. 5, 26852700.
- [34] M. Khovanov, Categorifications from planar diagrammatics. *Jpn. J. Math.* 5 (2010), no. 2, 153181.
- [35] M. Khovanov, L. Rozansky, Virtual crossings, convolutions and a categorification of the $SO(2N)$ Kauffman polynomial. *J. Gkova Geom. Topol. GGT* 1 (2007), 116214.
- [36] M. Khovanov, L. Rozansky, Matrix factorizations and link homology. *Fund. Math.* 199 (2008), no. 1, 191.
- [37] M. Khovanov, L. Rozansky, Matrix factorizations and link homology. II. *Geom. Topol.* 12 (2008), no. 3, 13871425.
- [38] C. Manolescu, P. Ozsvth, S. Sarkar, A combinatorial description of knot Floer homology. *Ann. of Math. (2)* 169 (2009), no. 2, 633660.
- [39] C. Manolescu, P. Ozsvth, Z. Szab, D. Thurston, On combinatorial link Floer homology. *Geom. Topol.* 11 (2007), 23392412.
- [40] P. Ozsvth, J. Rasmussen, and Z. Szab, Odd Khovanov homology. Preprint (2007), <http://arxiv.org/abs/0710.4300>.
- [41] J. H. Przytycki, When the theories meet: Khovanov homology as Hochschild homology of links, *arXiv:math.GT/0509334*
- [42] A. N. Shumakovitch, Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots, *J. Knot Theory Ramifications* 16 (2007), no. 10, 14031412.
- [43] P. Seidel, I. A. Smith, A link invariant from the symplectic geometry of nilpotent slices. *Duke Math. J.* 134 (2006), no. 3, 453514.
- [44] J. Rasmussen, Khovanov homology and the slice genus, *math.GT/0402131*.

- [45] J-L. Loday, Cyclic Homology, Grund. Math. Wissen. Band 301, Springer-Verlag, Berlin, 1992 (second edition, 1998).
- [46] J. P. May, Simplicial Objects in Algebraic Topology, University of Chicago Press (1967).
- [47] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information, Cambridge University Press (2000).
- [48] O. Viro (2004), Khovanov homology, its definitions and ramifications, *Fund. Math.*, **184** (2004), pp. 317-342.
- [49] B. Webster, Khovanov-Rozansky homology via a canopolis formalism. *Algebr. Geom. Topol.* 7 (2007), 673699.
- [50] B. Webster, Knot invariants and higher representation theory I: diagrammatic and geometric categorification of tensor products. math.GT.arXiv:1001.2020.
- [51] B. Webster, Knot invariants and higher representation theory II: the categorification of quantum knot invariants. math.GT. arXiv:1005.4559
- [52] E. Witten, Quantum Field Theory and the Jones Polynomial, *Comm. in Math. Phys.*, **121**, (1989), 351-399.
- [53] E. Witten, Knot Invariants from Four-Dimensional Gauge Theory. Davide Gaiotto, Edward Witten. physics.hep-th. arXiv:1106.4789
- [54] E. Witten, Fivebranes and Knots. Edward Witten. physics.hep-th. arXiv:1101.3216